

Case Report

Clinical Outcomes of Vitamin D Repletion in a Patient with Insulin Resistance: A Case Report

Alexander A Minin*

Institute of Protein Research, Russian Academy of Sciences, 119334 Moscow, Russia

*Correspondence: Alexander A Minin, Institute of Protein Research, Russian Academy of Sciences, 119334 Moscow, Russia , E-mail: alexminin@gmail.com, DOI: 10.1042/TDOM.1.1.0002

Abstract

Vitamin D is a secosteroid hormone that has traditionally been recognized for its central role in calcium-phosphate homeostasis and skeletal health. In recent years, growing evidence has suggested that vitamin D may also influence a range of extra-skeletal functions, including immune modulation, inflammation regulation, and glucose metabolism. Of particular interest is its potential association with insulin sensitivity and the pathophysiology of insulin resistance.

Keyword: Insulin resistance, Vitamin D, High dose, Obesity, Treatment

Received date: Jan 12, 2025; Accepted date: Jan 21, 2025; Published date: Jan 31, 2025

Citation: Alexander A Minin (2025) Clinical Outcomes of Vitamin D Repletion in a Patient with Insulin Resistance: A Case Report. RMB, 1: 1

Copyright: © 2025, Alexander A Minin. All intellectual property rights, including copyrights, trademarks rights and database rights with respect to the information, texts, images, logos, photographs and illustrations on the website and with respect to the layout and design of the website are protected by intellectual property rights and belong to Publisher or entitled third parties. The reproduction or making available in any way or form of the contents of the website without prior written consent from Publisher is not allowed.

Introduction

Insulin resistance is a metabolic condition characterized by reduced responsiveness of peripheral tissues to circulating insulin, leading to compensatory hyperinsulinemia and an increased risk of developing type 2 diabetes mellitus and other metabolic disorders. Multiple observational studies have demonstrated an association between low serum vitamin D levels and increased prevalence of insulin resistance; however, interventional studies have yielded inconsistent outcomes regarding the therapeutic benefit of vitamin D supplementation.

Parameter	Baseline Findings
Serum 25-hydroxyvitamin D	Deficient
Insulin sensitivity	Reduced
Glycemic status	Consistent with insulin resistance
Body weight	Increased
Acanthosis nigricans	Present
Menstrual pattern	Irregular cycles

Table 1: Baseline clinical characteristics.

This case report describes the clinical course of an individual diagnosed with vitamin D deficiency and features suggestive of insulin resistance. The patient presented with biochemical evidence of reduced serum 25-hydroxyvitamin D levels along with metabolic abnormalities consistent with impaired insulin action. Clinical manifestations included features commonly associated with insulin resistance, such as acanthosis nigricans and weight gain, along with menstrual irregularities suggestive of endocrine imbalance [Table 1].

Clinical Parameter	Baseline	After 6 months
Serum vitamin D	Deficient	Within optimal range
Insulin sensitivity	Reduced	Improved
Glycemic control	Impaired	Improved
Body weight	Elevated	Reduced
Acanthosis nigricans	Present	Partial regression
Menstrual regularity	Irregular	More regular

Table 2: Clinical changes following six months of vitamin d supplementation.

A high-dose vitamin D supplementation regimen was initiated following baseline evaluation. The supplementation was administered over a six-month period, with periodic clinical and biochemical monitoring to assess metabolic response and safety outcomes. The primary objective of the intervention was to correct vitamin D deficiency and observe potential downstream effects on insulin sensitivity and associated clinical features.

Over the course of treatment, a progressive increase in serum vitamin D levels was observed, eventually reaching values within the optimal physiological range. This improvement in vitamin D status coincided with notable metabolic and clinical changes. The patient exhibited improved insulin sensitivity, as reflected by better glycemic control indicators and reduced clinical signs of insulin resistance. A gradual regression of acanthosis nigricans was also noted, suggesting improvement in underlying hyperinsulinemic state [Table 2].

In addition to metabolic improvements, the patient experienced a reduction in body weight over the follow-up period. This weight change may have contributed further to improved insulin responsiveness. Furthermore, menstrual cycles became more regular, indicating possible restoration of endocrine balance, potentially mediated through improved metabolic function and vitamin D-related hormonal interactions.

Mechanism	Proposed effect	Potential clinical outcome
Improved pancreatic β -cell function	Enhanced insulin secretion	Better glucose regulation
Increased insulin receptor expression	Improved peripheral insulin responsiveness	Reduced insulin resistance
Anti-inflammatory effects	Reduced chronic inflammation	Improved metabolic function
Regulation of adipokines	Improved adipose tissue metabolism	Better weight regulation
Hormonal modulation	Improved endocrine balance	More regular menstrual cycles

Table 3: Proposed biological mechanisms of vitamin d in insulin resistance.

The observed clinical improvements may be explained through several proposed biological mechanisms. Vitamin D is thought to influence insulin secretion through its action on pancreatic β -cells and may enhance insulin sensitivity by modulating inflammatory pathways and upregulating insulin receptor expression in peripheral tissues. It may also play a role in regulating adipokine secretion, thereby influencing energy metabolism and adiposity [Table 3].

Despite these favorable observations, it is important to acknowledge that current evidence regarding vitamin D supplementation and insulin resistance remains inconclusive. While some studies suggest beneficial metabolic effects, others have reported

minimal or no significant improvement in insulin sensitivity following supplementation. This variability may be attributed to differences in baseline vitamin D status, dosage regimens, study duration, genetic factors, and heterogeneity in study populations.

Conclusion

Therefore, although this case supports a potential association between vitamin D repletion and metabolic improvement, it does not establish a definitive causal relationship. The findings should be interpreted cautiously, and high-quality randomized controlled trials with larger sample sizes are necessary to better define the therapeutic role of vitamin D in the management of insulin resistance and related metabolic disorders.

REFERENCES

- 1) Morris RL, PJ Hollenbeck. The regulation of bidirectional mitochondrial transport is coordinated with axonal outgrowth. *J Cell Sci.* 1993, 104:917-927.
- 2) Summerhayes IC, Wong D, Chen LB. Effect of microtubules and intermediate filaments on mitochondrial distribution. *J Cell Sci.* 1983, 61:87-105.
- 3) Jung D, Filliol D, Miehle M, Rendon A. Interaction of brain mitochondria with microtubules reconstituted from brain tubulin and MAP 2 or TAU. *Cell Motil. Cytoskel.* 1993, 24:245-255.
- 4) Leterrier JF, Rusakov DA, Nelson BD, Linden M. Interactions between brain mitochondria and cytoskeleton: Evidence for specialized outer membrane domains involved in the association of cytoskeleton-associated proteins to mitochondria in situ and in vitro. *Micros Res Tech.* 1994, 27:233-261.
- 5) Shea TB, Beermann ML, Fischer I. Transient requirement for vimentin in neuritogenesis: Intracellular delivery of anti-vimentin antibodies and antisense oligonucleotides inhibit neurite initiation but not elongation of existing neurites in neuroblastoma. *J Neurosci Res.* 1993, 36:66-76.
- 6) Menet V, Prieto M, Privat A, Ribotta MGY. Axonal plasticity and functional recovery after spinal cord injury in mice deficient in both glial fibrillary acidic protein and vimentin genes. *Proceedings of the National Academy of Sciences.* 2003, 100:8999-9004.
- 7) Boyne LJ, Fischer I, Shea TB. Role of vimentin in early stages of neuritogenesis in cultured hippocampal neurons. *Int J Dev Neurosci.* 1996, 14:739-748.
- 8) Parfenteva OI, Dayal AA, Gebreselase BA, Gyoeva FK, Alieva IB, et al. The Role of vimentin in neuritogenesis. *Russ J Dev Biol.* 2025, 56:40-46.
- 9) Vakhrusheva A, Endzhievskaya S, Zhuikov V, Nekrasova T, Parshina E, et al. The role of vimentin in directional migration of rat fibroblasts. *Cytoskeleton.* 2019, 76:467-476.
- 10) Nekrasova OE, Mendez MG, Chernoiivanenko IS, Tyurin-Kuzmin PA, Kuczmariski ER, et al. Vimentin intermediate filaments modulate the motility of mitochondria. *Mol Biol Cell.* 2011, 22:2282-2289.
- 11) Dayal AA, Medvedeva NV, Minin AA. N-Terminal fragment of vimentin is responsible for binding of mitochondria in vitro. *Membr Cell Biol. (Biochemistry (Moscow) Supplement Series A).* 2022, 5:21-28.
- 12) Chernoiivanenko IS, Matveeva EA, Gelfand VI, Goldman RD, Minin A. Mitochondrial membrane potential is regulated by vimentin intermediate filaments. *FASEB J.* 2014, 29:820-827.
- 13) Winter L, Abrahamsberg C, Wiche G. Plectin isoform 1b mediates mitochondrion-intermediate filament network linkage and controls organelle shape. *J Cell Biol.* 2008, 181:903-911.
- 14) Matveeva EA, Chernoiivanenko IS, Minin AA: Vimentin intermediate filaments protect mitochondria from oxidative stress. *Membr Cell Biol. (Biochemistry (Moscow) Supplement Series A).* 2010, 4:321-331.
- 15) Dayal AA, Parfenteva OI, Wang H, Gebreselase BA, Gyoeva FK, et al. Vimentin intermediate filaments maintain membrane potential of mitochondria in growing neurites. *Biol.* 2024, 13:995-1004.
- 16) Ivaska J, Vuoriluoto K, Huovinen T, Izawa I, Inagaki M, et al. PKC-mediated phosphorylation of vimentin controls integrin recycling and motility. *EMBO.* 2005, 24:3834-3845.
- 17) Burgstaller G, Gregor, M, Winter L, Wiche G. Keeping the vimentin network under control: cell-matrix adhesion-associated plectin 1f affects cell shape and polarity of fibroblasts. *Mol Biol Cell.* 2010, 21:3362-3375.
- 18) Strouhalova K, Přečková M, Gandalovičová A, Brábek J, Gregor M, et al. Vimentin intermediate filaments as potential target for cancer treatment. *Cancers.* 2020,184.