

Research Article

An Anti-SARS-CoV-2 Formula Consisting Volatile Oil from Traditional Chinese Medicine (TCM) Through Inhibiting Multiple Targets In Vitro

Huan Zhang¹, Wuyan Guo², Jing Shen³, Bo Zhang^{4*} and Xiaorui Song⁵

¹Key Laboratory of Industrial Fermentation Microbiology, Ministry of Education, College of Bioengineering, Tianjin, China

²Department of TCM Network Pharmacology, Tianjin Key Laboratory of Early Druggability Evaluation of Innovative Drugs, Tianjin, China

³Asymchem Life Science Co., Ltd. Tianjin, 300457, China

⁴Department of Automation, Institute for TCM-X, Beijing, China

⁵Department of Infectious Diseases, Henan International Joint Laboratory of Children's Infectious Diseases, Henan, China

***Correspondence:** Bo Zhang, Department of Automation, Institute for TCM-X, Beijing, China

Abstract

At present, coronavirus severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) and its variant virus are prevalent all over the world, taking a major toll on human lives worldwide. Oral mucosa and saliva are the high-risk routes of transmission. Inactivation of the virus in the mouth is one of the important strategies to reduce the source infectivity of the virus. However, the current preparations for inactivating oral virus are mainly aimed at Angiotensin Converting Enzyme2 (ACE2) protein, and its effect needs to be improved. In this study, through literature mining, multi-target screening and other methods, we screened and optimized the volatile oil formula (*Artemisiae argyi* folium, *Chrysanthemum morifolium*, *Trollii chinensis* flos, *Lonicerae japonica* flos) from the volatile oil of Traditional Chinese Medicine (TCM) to inhibit the invasion and replication of SARS-CoV-2. A test using pseudotype virus with S glycoprotein confirmed that this formula could effectively bind to the S glycoprotein to prevent SARS-CoV-2 host cell entry. Molecular docking experiments showed that the multiple molecules in the formula might weakly bind to these targets including ACE2, cathepsin L, furin, M^{pro} or 3cl. In all, the volatile oil formula designed here offers a general affordable strategy to protect patients from SARS-CoV-2 infections through debulking or minimizing transmission to others.

Keywords: SARS-CoV-2, Formula, Traditional Chinese Medicine (TCM), Multiple targets, Volatile oil

Received date: Apr 04, 2023; **Accepted date:** Apr 12, 2023; **Published date:** Apr 28, 2023

Citation: Zhang H, Guo W, Shen J, Zhang B, Song X (2024) An Anti-SARS-CoV-2 Formula Consisting Volatile Oil from Traditional Chinese Medicine (TCM) Through Inhibiting Multiple Targets *In Vitro*. J Infect Dis Ther. 2:1.

Copyright: © 2023, Zhang H, et al. All intellectual property rights, including copyrights, trademarks rights and database rights with respect to the information, texts, images, logos, photographs and illustrations on the website and with respect to the layout and design of the website are protected by intellectual property rights and belong to Publisher or entitled third parties. The reproduction or making available in any way or form of the contents of the website without prior written consent from Publisher is not allowed.

INTRODUCTION

Globally, the severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) outbreak has caused millions of infections and tens of thousands of deaths. The spread of SARS-CoV-2 occurs both through droplets and aerosols, and is strongly linked to indoor exposures of infected individuals, whether they are symptomatic or not [1]. In saliva, high levels of SARS-CoV-2 virus are often detected [2]. Both asymptomatic and symptomatic COVID-19 patients have high SARS-CoV-2 viral loads in saliva [3,4]. In fact, COVID-19 symptoms include the loss of taste and smell, and viral replication occurs in salivary glands and oral mucous membranes [5]. Therefore, Virus inactivation within the oral cavity may be an effective strategy to reduce the spread of SARS-CoV-2 through the oral mucous membranes and saliva.

A better understanding of how SARS-CoV-2 take over the host during infection is essential for constructing therapeutic strategies. In fact, SARS-CoV-2 replication relies on many proteins or enzymes from the virus itself and the host cell [6]. In principle, these proteins and enzymes represent potential therapeutic targets. Coronaviruses start their life cycles when their S proteins bind to Angiotensin-Converting Enzyme 2 (ACE2) on host cells' surface membranes [7]. According to a

previous description, the pre-fusion conformation is initiated by the proteolytic cleavage of the S1/S2 site, catalyzed by serine-proteases like furin or cathepsin L in correspondence of the (R/K)-(2X)n-(R/K) motif of S [8,9]. As a result of furin cleavage of the S1 (SPRRARY S) site of SARS-CoV-2, spike glycoprotein activation takes place, allowing the virus to enter the host. It is a polybasic cleavage site found in SARS-CoV-2, also known as a multibasic cleavage site. In COVID-19 patients, Cathepsin-L is thought to play a critical role since its levels are elevated after SARS-CoV-2 infection and positively correlate with disease severity [10]. Functionally, SARS-CoV-2 infection is enhanced by cathepsin-L by cleaving furin-primed SARS-CoV-2 S protein into smaller fragments and promoting cell-cell fusion [10,11]. Subsequently, the coronavirus main proteinase (3CL^{PRO}) of the virus cleaves viral replicase polyprotein into effector proteins [12]. Considering their undeniable role in the entry and replication of viruses, it may be possible to inhibit or modulate these four proteins (Angiotensin- Converting Enzyme (ACE2), furin, cathepsin L and 3CL^{PRO}) to develop effective antiviral therapies.

Currently, the oral preparation used to inactive SARS-CoV-2 are mainly aiming to act on the ACE2 target, which plays a major role in the invasion of host. Daniell et al., developed a chewing gum with converting enzyme 2 proteins to decrease oral virus transmission and infection through debulking SARS-CoV-2 in saliva [13]. Another study demonstrated newly synthesized polypeptides inhibits eliminating the invading virus by competing with SARS-CoV-2 for the ACE2 receptors [14]. However, these Oral preparation have been exclusively focused on one single protein target. Based on the fact that viruses go through various stages of their life cycle, an effective strategy involves targeting multiple viral proteins it enables the potential preparation to disrupt the viral infection and replication process in different stages as a result, the effect of killing viruses is maximized [15].

Recent studies showed that Traditional Chinese Medicine (TCM) could acts on multiple targets and multiple pathological processes, which provided us a valuable resource [16]. Traditional Chinese medicine volatile oil rich in ketones, terpenes, aldehydes, alcohols and other active substances is the product of plant metabolism extracted from plants by specific methods [17]. It is widely used in development of drugs, antibacterial agents, antiviral agents and other products due to its activities of enhancing immunity, bacteriostasis and antiviral [18,19]. Moreover, it has aromatic components, and the raw materials are simple and easy to obtain. Compared with traditional chemical drug products, volatile oil can possess more efficient antibacterial activity and are not easy to produce resistance limitation, with lower cytotoxicity [19,20]. Thus, it is feasible to develop safe and environment-friendly anti-SARS- COVID-2 oral preparation from the volatile oil of traditional Chinese medicine.

In this research, we screened from the volatile oils of Traditional Chinese Medicine based on literature mining and multiple target screening and obtain an effective formula. Then, the effect of the formula was investigated by *in vitro* assay. Finally, we use molecular docking to preliminarily explore the mechanism of this compound against SARS-CoV-2.

MATERIALS AND METHODS

Reagents and cell culture

HEK293T-overexpressing ACE2 cell were purchased from GENEWIZ (Jiangsu, China) and grown in Dulbecco's Modified Eagle Medium (DMEM) (gibco, thermo fisher scientific, waltham, MA, USA) containing 10% fetal bovine serum (Gibco; hereinafter referred to as complete medium), 100 U/mL penicillin, 100 µg/mL streptomycin and 1 µg/mL puromycin and incubated at 37° C in an atmosphere containing 5% CO₂/95% air.

Literature mining

All Raw data is uploaded as supplementary file. All data generated or analyzed during this study are included in this published article and its supplementary information files.

"Traditional Chinese medicine", "volatile oil" and "antiviral" were used as key words for a comprehensive search of both english and chinese-language literature in PubMed, and CNKI (Chinese National Knowledge Infrastructure) databases from May 2000 to May 2022. For overlapping or republished studies, only the most recent published papers were included. There are 214 references and 62 kinds of single Traditional Chinese Medicine volatile oils are retrieved. Finally, 9 kinds of Traditional Chinese Medicine volatile oils was obtained according to the *in vitro* antiviral mechanism.

Extraction of volatile oil by hydrodistillation

5 kgs of 9 kinds TCM were purchased from anguo kanghua Traditional Chinese Medicine sales Co. Ltd. (Hebei, China).

Each of above TCM was subjected to hydrodistillation for 2 h, to extract volatile oil, and the volatile oil was then kept in a glass bottle after cooling down to the room temperature. The residual water was then removed, using anhydrous sodium

sulfate. The yield of essential oil was calculated by using the following formula:

$$\%Yield = \left(\frac{a}{b} \right) \times 100$$

Where, a is a volume of the volatile oil, and b is the weight of fresh plant materials used in the hydrodistillation. The essential oil was stored at 4°C, in a light-protected container, until further use.

Cathepsin L activity assays

The cathepsin L activity screening assay kit (Biovision, California, USA) was used to determine the inhibitory effect of selected volatile oils on the activity of isolated cathepsin L enzyme according to the manufacturer's protocol. Selected volatile oils at 0.5 mg/mL concentrations were added to cathepsin L (0.2 mU/μL) and the reaction mix was incubated for 15 min at RT. Positive control consisted of cathepsin L alone, whereas negative control consisted of cathepsin L and inhibitor (10 μM) of cathepsin L. Cathepsin L substrate was added to each well. The fluorescence was measured at Ex/Em=400/505 nm wavelength using a spectrofluorometer (thermofisher, Massachusetts, USA) in a kinetic mode for 30 min at 37°C. Control was 0.5% ethanol. Results are expressed as a percentage of volatile oil-free control (mean ± SD, n=6).

Angiotensin Converting Enzyme2 (ACE2) activity assays

To determine the inhibitory effect of selected volatile oils on the activity of recombinant hACE2 protein, an ACE2 activity screening assay kit (Beyotime, Shanghai, China) was used according to the manufacturer's protocol. Briefly, the volatile oils at 0.5 mg/mL concentrations were added and the reaction mix was incubated for 45 min at RT. The positive control was a sample containing ACE2 enzyme and ACE2 enzyme inhibitor MLN-4760 (10 nm). The fluorescence was measured at Ex/Em=325/393 nm wavelength using a spectrofluorometer (thermofisher, Massachusetts, USA). 100% enzyme activity control was 0.5% ethanol. Results are expressed as a percentage of volatile oil-free control (mean ± SD, n=6).

Furin activity assays

To examine the effect of various selected volatile oils on furin activity, a furin Activity Screening Assay Kit (Biovision, California, USA) was used according to the manufacturer's protocol. Briefly, the volatile oils at 0.5 mg/mL concentrations were added. furin substrate was added to each well. The positive control was a sample containing furin enzyme and furin inhibitor (10 nm). The fluorescence was measured at Ex/Em=360/460 nm wavelength using a spectrofluorometer (thermofisher, Massachusetts, USA) in a kinetic mode for 1 h at 37°C. Results are expressed as a percentage of volatile oil-free control (mean ± SD, n=6).

M^{pro} or 3cl activity assays

To determine the inhibitory effect of selected volatile oils on the activity of M^{pro} or 3cl protein, an M^{pro} or 3cl activity screening assay kit (Beyotime, Shanghai, China) was used according to the manufacturer's protocol. Briefly, the volatile oils at 0.5 mg/mL concentrations were added and the reaction mix was incubated for 45 min at Respiratory Therapist (RT). The positive control was a sample containing M^{pro} or 3cl enzyme and M^{pro} or 3cl enzyme inhibitor Ebselen (2 μM). The fluorescence was measured at Ex/Em=325/393 nm wavelength using a spectrofluorometer (thermofisher, Massachusetts, USA). 100% enzyme activity control was 0.5% ethanol. Results are expressed as a percentage of volatile oil-free control (mean ± SD, n=6).

Viability assay

In order to assess cell viability, we used the CCK-8 assay. Briefly, HEK-293T/ACE2 cells were seeded into a 96-well plate at a cell density of 6,000 per well, and allowed to adhere for 24 h, followed by treatment with different concentrations of selected volatile oils for up to 48 h. Afterwards, complete growth medium was replaced with a fresh one substituted with 10 μL CCK-8, followed by 3 hours at 37°C incubation. The absorbance was measured at 450 nm using a spectrophotometer (Molecular Devices, San Jose, CA). Results are expressed as a percentage of volatile oils-free control (mean ± SD, n=6).

Pseudovirus preparation and neutralization assay

For pseudovirus neutralization assay, 100 Tissue Culture Infectious Dose (TCID50) per well pseudovirus in medium without FBS was incubated with volatile oils combination for 30 min at room temperature. Then the mixtures (100 μL) were used to infect HEK- 293T/ACE2 cells rinsed with PBS. After 5 h, 100 μL medium with 5% FBS was added. After incubation for another 48 h, After 48 h, cells were lysed using Firefly Luciferase Assay Kit (Genescript, catalog no. L00877C), and relative

light units were measured using a 96 Thermo Microplate luminometer. Percent neutralization was calculated using GraphPad Prism 8.

Molecular docking

Ligand preparation: For molecular docking studies, The structures of the all ligands were retrieved from European Molecular Biology Laboratory (EMBL)- European Bioinformatics Institute (EBI) (www.ebi.ac.uk/chebi/advancedSearchFT.do), in a three-dimensional Structure File (SDF) format, and furthermore, the structure was refined using the ligprep module in schrodinger's maestro (v 12.8). The OPLS4 force field was applied, and 32 different states of stereoisomerism were derived (Schrödinge release 2021-2: ligprep, Schrödinge, LLC, New York, NY, 2021).

Protein preparation: We need to evaluate the furin (5JXH), M^{PRO} (6W63), cathepsin L (3OF9) and anti ACE2 (1R42) activity against different ingredients in the each volatile oils computationally, respectively. The three-dimensional structure of proteins was retrieved from the database of Protein Data Bank (PDB). The X-ray crystallographic structures were imported into Maestro using the protein preparation wizard, and this module helps to solve the missing hydrogen bonds, create the disulfide bonds, and optimize (Schrödinge Release 2021-2: Protein Preparation Wizard; Epik, Schrödinge, LLC, New York, NY, 2021; Impact, Schrödinge, LLC, New York, NY; Prime, Schrödinge, LLC, New York, NY, 2021).

The molecular docking was performed using the Glide package (ligand docking) in the Schrodinger suite. The standard precision docking method was applied and performed postdocking minimization and analyzed the results in pose viewer Schrödinge Release 2021-2: Glide, Schrödinge, LLC, New York, NY, 2021

TCM	Scientific name	Refractive index	pH	Density
Shi xiang ru	<i>Mosiae Herba</i>	1.4718	4.4	0.9012
Ma ye qian li guang	<i>Senecionis scandentis hebra</i>	1.4716	4.4	0.9555
Yu xing cao	<i>Houttuyniae herba</i>	1.3408	4.4	0.8756
Hong feng	<i>Acer palmatum 'atropurpureum'</i>	1.339	4.4	0.9227
Ai ye	<i>Artemisiae argyi folium</i>	1.3334	4.4	0.9946
Jin lian hua	<i>Lonicerae japonica flos</i>	1.3343	4.4	0.8768
Hang bai ju	White <i>Chrysanthemum indicum</i>	1.3338	4.1	0.96
Ye ju hua	<i>Chrysanthemi indici flos</i>	1.3361	4.4	0.8346
Jin yin hua	<i>Trollius chinensis bunge</i>	1.4679	4.5	0.872

Table 1: Physicochemical characteristics of 9 volatile oils.

Statistical analysis

All data in experiments were analyzed using Prism 8.0 software and are presented as mean \pm SEM. Statistical significance was assessed by one-way Analysis of Variance (ANOVA). P value <0.05 indicates significance.

RESULTS AND DISCUSSION

Antiviral research has been an active field in recent decades. To date, at present, many Traditional Chinese Medicines have been reported to have antiviral effects. In the long history of Chinese medicine, some TCMs volatile oil have been reported to have antiviral benefits. Thus, TCMs volatile oil are great candidates for screening and verifying their antiviral properties. To choose TCM volatile oil candidates for validation, we searched the National Pubmed as well as China National Knowledge Infrastructure (CNKI) for antiviral TCMs volatile oil. The mechanism of Traditional Chinese Medicine can be divided into direct inhibition and indirect inhibition. The direct inhibition is mainly to block a certain link in the process of virus reproduction to defeat pathogen propagation; Indirect inhibition is to stimulate and mobilize the immune defense system of the body to play an antiviral role [21]. Since we only consider the direct antiviral effect, that is, to inhibit the virus invasion and replication in the host, we finally get 9 Chinese herbal volatile oils through screening, as shown in Table 1.

Four volatile oils inhibit the activity of different targets of SARS-CoV-2, respectively

As the life cycle of SARS-CoV-2 is related to multiple protein targets, as shown in Figure 1 and 2, in order to better prevent SARS-CoV-2 from invading and replicating, we screened volatile oils that can inhibit these four targets, including ACE2, furin, cathepsin L and M^{pro} or 3cl. The angiotensin-converting enzyme II (ACE2) has been recognized as the host receptor of severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2). There is a strong interaction between the receptor binding domain of the SARS-CoV-2 S-protein and ACE2, in this way, ACE2 may provide an effective drug target for preventing COVID-19 from invading host cells [22]. The screening results are shown in Figure 3A. The volatile oil of Hang bai ju (White *Chrysanthemum indicum*) is the best one to target ACE2.

Furin converts synthesized proteins into biologically active forms by cleaving a specific section [23]. It is a serine endoprotease that cleaves precursor protein processing sites in a calcium-dependent manner. Furin cleaves S1 (SPRRARY S) site of SARS-CoV-2, direct emergence in spike glycoprotein activation for viral entry into the host system [24]. The screening results are shown in Figure 3B. Among the volatile oils of Traditional Chinese Medicine targeting furin protease, the volatile oil of Ai ye (*Artemisiae argyi* folium) has the best inhibitory effect.

Cathepsin-L plays a critical role in COVID-19 patients, as evidenced by high circulating levels after SARS-CoV-2 infections, and a positive correlation with disease severity [25]. Functionally, It helps promote SARS-CoV-2 infection by fragmenting furin-primed SARS-CoV-2 S proteins and facilitating cell-to-cell fusion by cleaving them into smaller fragments [9,11]. Results as shown in Figure 3C, the inhibitory effect of honeysuckle volatile oil on cathepsin L was the best.

M^{pro} is highly conserved among different CoVs, and as far as we known, It has no human homolog, and shows a low mutation rate, making it ideal for the development of multi-viral drugs that can interfere with different varieties of SARS-CoV-2 and other coronaviruses' vital cycles [26,27]. The screening results are shown in Figure 3D. Among the volatile oils of Traditional Chinese Medicine targeting the main protease, the volatile oil of Jin lian hua (*Trollius chinensis*) has the best inhibitory effect.

Neutralization assay with SARS-CoV-2 spike-pseudotyped lentiviral particles

For maximum antiviral effect, we design a combination containing 0.25 mg/mL. The anticoronavirus activity of the formula was further explored by pseudovirus neutralization test. Pseudoviruses containing the viral spike protein and expressing the luciferase reporter gene were used to infect Human Embryonic Kidney Cells. (HEK293) expressing human ACE2. The formula at the indicated concentration was incubated at room temperature for 90 minutes with spike glycoprotein pseudotyped viruses expressing luciferase expressed by the SARS-CoV-2 virus. Then, virus-containing the formula was incubated with ACE2-expressing HEK293 cells for 72 h, and viral infectivity was measured via luciferase. The incubation of pseudotyped lentivirus with the four kind volatile oil and the formula significantly reduced luciferase activities, and at 1 mg/ml of the highest inhibition was observed (Figure 4), confirming the ability of this formula to block entry of lentivirus spike protein into HEK293 cells either by direct inhibiting ACE2 or cathepsin L or furin or M^{pro} or 3cl protein on HEK293 cells. In addition, the effects of the formula on the cell viability were also evaluated, and it demonstrated no cytotoxicity (Figure 5)

The volatile oil in this article were rationally designed to target multiple targets either on viral entry or replication. The volatile oil in this article is a multiple target targeting COVID-19, This concept is consistent with the idea of drug discovery targeting COVID-19 multi protein reported previously [15]. The definition of a drug target is important to the success of drug discovery [28]. An antiviral agent targeting multiple viral proteins is a novel concept. Similar to drug cocktail screening, drug combination screening involves selecting drug candidates that target and block different stages of the virus' life cycle. Contrary to the majority of existing repurposed drug research, this work focuses on multiple targets at once. Thus, our approach has the potential to attack the virus from different angles if successful.

Exploring the antiviral mechanism of formula based on molecular docking

The antiviral mechanisms of four each volatile oil could preliminarily obtained through the literature. It is reported that the volatile oil of Ai ye (*Artemisiae argyi* folium) has inhibitory effect on herpes zoster virus [29], Respiratory Syncytial Virus (RSV), Influenza Virus (IFV) and Hepatitis B Virus (HBV) [30,31]. Artemisol and other components can inhibit bacteria and viruses [31]. Linalool, geraniol and a-terpineol from the volatile oil of Jin yin hua (*Trollius chinensis* bunge) hua have antiviral effects [32]. Hai et al., reported that volatile compounds are one of the main active ingredients of Hang bai ju (White *Chrysanthemum indicum*), which have a significant anti influenza virus (H3N2) effect [33,34]. Zhao et al., found that the ethanol extract of Jin lian hua (*Trollius chinensis*) has obvious inhibitory effect on virus replication in chicken embryos, and

has direct killing effect on H1N1 virus PR8 strain [35].

We next asked what is the specific mechanism underlying anti- COVID19 efficacy of the formula. It is necessary to explicitly determine the main components of these volatile oils. According to reports in the literature, volatile oil of Hang bai ju (White *Chrysanthemum indicum*), Ai ye (*Artemisiae argyi* folium), Jin yin (*Trollius chinensis* bunge) hua and jin lian hua (*Trollius chinensis*) mainly contains components respectively, as shown in Tables 2-5. To investigate the interactions between component of volatile oil and different targets, molecular docking experiments were performed [36-39]. The component of the four kinds of essential oil were docked within the active site of four target using Extra Precision (XP) mode of Glide docking module of Maestro, respectively. Based on the docking scores, the compounds were ranked. PyMOL was used to visualize ligand-receptor interactions within the protein's active site. The results shown that isocyclocitral from Hang bai ju (White *Chrysanthemum indicum*), spathulenol from Ai ye (*Artemisiae argyi* folium), benzyl benzoate from Jin yin hua (*Trollius chinensis* bunge), and beta-ionone from Jin lian hua (*Trollius chinensis*) bind at the active site of ACE2, furin, cathepsinL and M^{Pro} or 3cl of SARS-CoV-2 with the dock score -4.007 kcal/mol, -4.995 kcal/mol, -6.146 kcal/mol and -4.927 kcal/mol, respectively (Tables 2-5 and Figure 6). Moreover, it also demonstrated that other components of each volatile oil could act on the corresponding target, even though the interaction between them is very weak.

There are only five components of the volatile oil from Jin yin hua (*Trollius chinensis* bunge), whose docking score with absolute values were greater than 5, indicating that the binding ability of these volatile oil monomers to the target was weak. As is well known to all, herbs produce their efficacy through the synergistic effects of multi- ingredients, multi-targets, and multi-pathways [40,41]. Although the results show that the absolute value of the docking score between these volatile oil monomers and the target is relatively small, these volatile oils with multi-component synergizing on the common target can also achieve the effect of acting on the target.

ID	Name	Glide ligand efficiency	Glide gscore	Glide energy
CHEBI:171934	Isocyclocitral	-0.364	-4.007	-17.951
CHEBI:89773	1,2-Dihydro-1,1,6-trimethylnaphthalene	-0.277	-3.596	-17.596
CHEBI:167345	Cedrene	-0.238	-3.564	-19.948
CHEBI:167347	gamma-Selinene	-0.226	-3.383	-15.157
CHEBI:10357	(-)-beta-Caryophyllene	-0.187	-2.807	-9.286
CHEBI:180521	D6-Ambrettolide	-0.154	-2.763	-12.213
CHEBI:63190	(+)-beta-Caryophyllene	-0.179	-2.68	-10.828
CHEBI:62224	(-)-7-epi-alpha-Selinene	-0.175	-2.629	-15.377
CHEBI:62757	alpha-Curcumene	-0.169	-2.529	-17.887
CHEBI:64361	beta-Sesquiphellandrene	-0.164	-2.465	-15.725
CHEBI:84851	Ethyl linolenate	-0.085	-1.862	-30.722
CHEBI:31572	Ethyl linoleate	-0.071	-1.552	-25.76
CHEBI:34687	Dibutyl phthalate	-0.034	-0.681	-27.935
CHEBI:32936	Tetracosane	0.001	0.013	-26.007
CHEBI:46050	Docosane	0.002	0.037	-22.23
CHEBI:17327	Phytol	0.029	0.611	-25.452
CHEBI:39238	(Z,E)-alpha-Farnesene	0.049	0.742	-17.625
CHEBI:17351	Linoleic acid	0.071	1.42	-22.532
CHEBI:69080	Methyl linoleate	0.072	1.515	-25.416
CHEBI:10418	trans-beta-Farnesene	0.111	1.669	-18.428
CHEBI:30813	Decanoic acid	0.161	1.927	-14.954
CHEBI:15756	Hexadecanoic acid	0.133	2.394	-21.115
CHEBI:43619	Icosane	0.154	3.077	-22.779
CHEBI:16148	Heptadecane	0.189	3.221	-20.402
CHEBI:32927	Nonadecane	0.171	3.249	-21.935
CHEBI:32926	Octadecane	0.189	3.405	-20.733
CHEBI:132293	2-Methylicosane	0.162	3.407	-21.296

Table 2: Molecular docking results of the essential oil from hang bai ju (White *Chrysanthemum indicum*) with ACE2.

ID	Name	Glide ligand efficiency	Glide gscore	Glide energy
CHEBI:132824	Spathulenol	-0.312	-4.995	-29.469
CHEBI:167328	2-p-Menthene-1-ol	-0.435	-4.781	-22.01
CHEBI:176780	Dihydroactinidiolide	-0.365	-4.739	-21.911
CHEBI:50040	Thujone	-0.429	-4.715	-18.481
CHEBI:4609	Dimethyl phthalate	-0.322	-4.514	-27.789
CHEBI:17854	Cyclohexanone	-0.636	-4.454	-18.001
CHEBI:22469	alpha-Terpineol	-0.405	-4.454	-26.024
CHEBI:173721	Davanone	-0.262	-4.453	-29.093
CHEBI:27961	1,8-cineole	-0.39	-4.29	-19.998
CHEBI:3573	Chamazulene	-0.301	-4.209	-22.475
CHEBI:132905	alpha-Cadinol	-0.256	-4.094	-23.184
CHEBI:28093	Borneol	-0.372	-4.088	-19.503
CHEBI:132899	beta-Terpineol	-0.371	-4.078	-26.049
CHEBI:29069	Phthalic acid	-0.336	-6.432	-30.927
CHEBI:36773	Camphor	-0.366	-4.03	-18.778
CHEBI:156228	Viridilorol	-0.251	-4.017	-23.82
CHEBI:15397	(S)-Camphor	-0.364	-4.003	-18.068
CHEBI:49045	Germacrene D	-0.26	-3.893	-22.31
CHEBI:10577	gamma-Terpinene	-0.368	-3.684	-18.022

CHEBI:61699	(-)-alpha-Gurjunene	-0.246	-3.683	-25.021
CHEBI:63709	Bicyclogermacrene	-0.24	-3.595	-24.5
CHEBI:50025	beta-Pinene	-0.331	-3.31	-15.545
CHEBI:17327	Phytol	-0.072	-1.503	-39.271
CHEBI:145715	Hex-3-en-1-ol	-0.158	-1.108	-17.563
CHEBI:185431	Pentane-1,5-diol	-0.143	-1.001	-23.626
CHEBI:39242	cis-beta-Farnesene	0.05	0.743	-25.779
CHEBI:17302	Pentadecanal	0.136	2.173	-30.775

Table 3: Molecular docking results of the essential oil from Ai ye (*Artemisiae argyi* folium) with furin.

ID	Name	Glide ligand efficiency	Glide gscore	Glide energy
CHEBI:41237	Benzyl benzoate	-0.384	-6.146	-32.778
CHEBI:28266	Fluorene	-0.454	-5.901	-24.18
CHEBI:22469	Alpha-terpineol	-0.535	-5.89	-25.072
CHEBI:28851	Phenanthrene	-0.416	-5.83	-24.734
CHEBI:34247	2,6-di-tert-butyl-4-methylphenol	-0.342	-5.472	-23.883
CHEBI:32319	Alpha-ionone	-0.35	-4.906	-23.664
CHEBI:84851	Ethyl linolenate	-0.17	-3.745	-36.322
CHEBI:32936	Tetracosane	-0.15	-3.595	-35.184
CHEBI:17447	Geraniol	-0.325	-3.575	-23.114
CHEBI:32941	Heptacosane	-0.114	-3.082	-39.933
CHEBI:32938	Pentacosane	-0.106	-2.659	-34.403
CHEBI:17580	Linalool	-0.236	-2.598	-21.637
CHEBI:149547	(2E,4E)-deca-2,4-dienal	-0.233	-2.568	-24.047

CHEBI:46050	Docosane	-0.1	-2.211	-35.118
CHEBI:34687	Dibutyl phthalate	-0.098	-1.956	-39.578
CHEBI:67252	Farnesyl acetone	-0.097	-1.85	-32.469
CHEBI:17351	Linoleic acid	-0.092	-1.846	-38.83
CHEBI:177721	Methyl 9-oxononanoate	-0.073	-0.946	-29.415
CHEBI:17327	Phytol	-0.033	-0.683	-32.077
CHEBI:42504	Pentadecanoic acid	-0.039	-0.664	-33.772
CHEBI:15756	Hexadecanoic acid	-0.024	-0.442	-36
CHEBI:28842	Octadecanoic acid	-0.02	-0.397	-39.091
CHEBI:30805	Dodecanoic acid	-0.028	-0.396	-29.905
CHEBI:30813	Decanoic acid	-0.033	-0.394	-28.055
CHEBI:69187	Methyl palmitate	-0.015	-0.277	-33.707
CHEBI:69080	Methyl linoleate	-0.011	-0.224	-35.128
CHEBI:28875	Tetradecanoic acid	0	-0.004	-30.723
CHEBI:69188	Methyl stearate	0.006	0.136	-36.556
CHEBI:84156	Methyl palmitoleate	0.016	0.312	-33.112
CHEBI:84067	Tetradecanal	0.064	0.956	-29.986
CHEBI:45296	Hexadecane	0.111	1.774	-28.734
CHEBI:32927	Nonadecane	0.099	1.884	-30.142

Table 4: Molecular docking results of the essential oil from Jin yin hua (*Trollius chinensis* bunge) with Cathepsin L.

ID	Name	Glide ligand efficiency	Glide gscore	Glide energy
CHEBI:32325	Beta-ionone	-0.352	-4.927	-27.773
CHEBI:67206	Geranyl acetone	-0.325	-4.554	-27.44
CHEBI:42438	2-methoxy-4-vinylphenol	-0.402	-4.422	-23.155
CHEBI:67252	Farnesyl acetone	-0.212	-4.035	-29.67
CHEBI:17351	Linoleic acid	-0.192	-3.843	-32.227
CHEBI:16196	Oleic acid	-0.187	-3.741	-35.214
CHEBI:46050	Docosane	-0.166	-3.663	-31.617
CHEBI:17580	Linalool	-0.329	-3.614	-22.597
CHEBI:28716	Palmitoleic acid	-0.183	-3.299	-28.459
CHEBI:79053	Diisobutyl phthalate	-0.164	-3.287	-31.689
CHEBI:69080	Methyl linoleate	-0.156	-3.276	-32.755
CHEBI:17327	Phytol	-0.147	-3.087	-26.609
CHEBI:28842	Octadecanoic acid	-0.154	-3.08	-30.355
CHEBI:27432	Alpha-linolenic acid	-0.144	-2.893	-31.987
CHEBI:29019	Nonanoic acid	-0.248	-2.735	-21.532
CHEBI:133634	Methyl linolenate	-0.13	-2.725	-35.695
CHEBI:42504	Pentadecanoic acid	-0.144	-2.444	-27.718
CHEBI:16148	Heptadecane	-0.144	-2.441	-29.375
CHEBI:28897	Pentadecane	-0.157	-2.355	-26.189
CHEBI:87494	Methyl laurate	-0.15	-2.253	-27.222
CHEBI:41808	decane	-0.221	-2.212	-17.68
CHEBI:28875	Tetradecanoic acid	-0.134	-2.15	-25.702
CHEBI:41253	Tetradecane	-0.132	-1.854	-27.426
CHEBI:35998	Tridecane	-0.123	-1.599	-26.364

CONCLUSION

In this work, we present a formula consisting volatile oil for anti- viral agents that interact with multiple target proteins of SARS-CoV-2. We first collected 9 antiviral volatile oils from the literature, Secondly, the volatile oil with the best inhibitory effect on the four targets was obtained by target screening. Then the formula consisting the four volatile oils has better effect of neutralizing pseudovirus compared with the effects of mono-agents. Overall, we found a volatile oil formula with potential inhibitory ability against 4 key proteins of SARS-CoV-2. The volatile oil can be used to develop oral or nasal preparations. A further study in animals is needed to determine if oral or nasal spray delivery is more effective as a preventative or curative measure. There are more than four targets related to the life cycle of novel coronavirus. In the future, we will further screen other targets to improve the efficacy of this formula.

AUTHOR CONTRIBUTION

Conceptualization was primarily conducted by Huan Zhang and Bo Zhang. Methodology and resources were developed by Huan Zhang, Wuyan Guo, Jing Shen and Xiaorui Song. Validation and visualization were carried out by Huan Zhang and Wuyan Guo. Formal analysis involved contributions from Huan Zhang, Wuyan Guo, and Bo Zhang. The original draft was prepared by HZ, with subsequent writing review and editing conducted by Huan Zhang, Wuyan Guo and Bo Zhang. Bo Zhang provided supervision throughout the project. All authors have read and agreed to the published version of the manuscript.

FUNDING

This research was funded by China postdoctoral science foundation (2020T130093ZX) and the open research funding (ERC202307).

DATA AVAILABILITY

The data used to support the findings of this study are available from the corresponding author upon request.

REFERENCES

- 1) Allen JG, Ibrahim AM (2021) Indoor air changes and potential implications for SARS-CoV-2 transmission. *Jama* 325:2112-2113.
- 2) Li Y, Ren B, Peng X, Hu T, Li J, et al. (2020) Saliva is a non-negligible factor in the spread of COVID-19. *Mol Oral Microbiol* 35:141-145.
- 3) Chan JF, Yip CC, To KK, Tang TH, Wong SC, et al. (2020) Improved molecular diagnosis of COVID-19 by the novel, highly sensitive and specific COVID-19- rdp/hel real-time reverse transcription-pcr assay validated *in vitro* and with clinical specimens. *J Clin Microbiol* 58:310-320.
- 4) Wölfel R, Corman VM, Guggemos W, Seilmaier M, Zange S, et al. (2020) Virological assessment of hospitalized patients with COVID-2019. *Nature* 581:465-469.
- 5) Huang N, Pérez P, Kato T, Mikami Y, Okuda K, et al. (2021) SARS-CoV-2 infection of the oral cavity and saliva. *Nat Med* 27:892-903.
- 6) Gil C, Ginex T, Maestro I, Nozal V, Barrado-Gil L, et al. (2020) COVID-19: Drug targets and potential treatments. *J Med Chem* 63:12359-12386.
- 7) Zhou P, Yang XL, Wang XG, Hu B, Zhang L, et al. (2020) A pneumonia outbreak associated with a new coronavirus of probable bat origin. *Nature* 579:270-273.
- 8) Coutard B, Valle C, De Lamballerie X, Canard B, Seidah NG, et al. (2020) The spike glycoprotein of the new coronavirus 2019-nCoV contains a furin-like cleavage site absent in CoV of the same clade. *Antiviral res* 176:104742.
- 9) Ou X, Liu Y, Lei X, Li P, Mi D, et al. (2020) Characterization of spike glycoprotein of SARS-CoV-2 on virus entry and its immune cross-reactivity with SARS-CoV. *Nat Commun* 11: 1620.
- 10) Zhao MM, Yang WL, Yang FY, Zhang L, Huang WJ (2021) Cathepsin L plays a key role in SARS-CoV-2 infection in humans and humanized mice and is a promising target for new drug development. *Signal Transduct Target Ther* 6:134-134.
- 11) Shang J, Wan Y, Luo C, Ye G, Geng Q, et al (2020). Cell entry mechanisms of SARS-CoV-2. *Proc Natl Acad Sci* 117:11727-11734.
- 12) Zhang L, Lin D, Sun X, Curth U, Drosten C, et al. (2020) Crystal structure of SARS-CoV-2 main protease provides a basis for design of improved alpha- ketoamide inhibitors. *Science*,368:409-412.
- 13) Daniell H, Nair SK, Esmaeili N, Wakade G, Shahid N, et al. (2021) Debulking SARS-CoV-2 in saliva using angiotensin converting enzyme 2 in chewing gum to decrease oral virus transmission and infection. *Mol Ther* 30:1966-78.
- 14) Zeng Z, Hu M, Wang X, Zheng B, Zeng C, et al. (2021) A disinfectant combination for preventing coronavirus 12:22-25.
- 15) Liang H, Zhao L, Gong X, Hu M, Wang H (2021) Virtual screening FDA approved drugs against multiple targets of SARS-CoV-2. *JCTS* 14:1123-1132.
- 16) Li L, Zhang L, Yang CC (2016) Multi-target strategy and experimental studies of traditional Chinese medicine for Alzheimer's disease therapy. *Curr Top Med Chem* 16:537-48.
- 17) Ren P, Fan N, Tian M, Qin Y (2018) Research progress on medical effects of essential oils. *J Tradit Chin Med* 33:2507-2511.
- 18) Birmpa A, Constantinou P, Dedes C, Bellou M, Sazakli E (2018) Antibacterial and antiviral effect of essential oils combined with non-thermal disinfection technologies for ready-to-eat romaine lettuce. *J Nutr Food Technol* 1:24-32.
- 19) Reichling J, Schnitzler P, Suschke U, Saller R (2009) Essential oils of aromatic plants with antibacterial, antifungal, antiviral, and cytotoxic properties an overview. *Complement Med Res* 16:79-90.

21) Reichling J, Schnitzler P, Suschke U, Saller R (2011) Antiviral effects of plant- derived essential oils and pure oil components. *Lipids and Essential Oils* 16:79- 90.

22) Li H, Jia X (2015) Research progress on the antiviral mechanism of traditional Chinese medicine. *J Tradit Chin Med* 6:82-85.

23) Yao, H (2022) Virtual screening of natural chemical databases to search for potential ACE2 inhibitors. *Molecules* 27:1740.

24) Kiefer MC, Tucker JE, Joh R, Landsberg KE, Saltman D, et al. (1991) Identification of a second human subtilisin-like protease gene in the fes/fps region of chromosome 15. *DNA cell biol* 10:757-769.

25) Vardhan S, Sahoo SK (2022) Virtual screening by targeting proteolytic sites of furin and TMPRSS2 to propose potential compounds obstructing the entry of SARS-CoV-2 virus into human host cells. *J Tradit Complement Med* 12:6-15.

26) Zhao MM, Yang WL, Yang FY, Zhang L, Huang WJ, et al. (2021) Cathepsin L plays a key role in SARS-CoV-2 infection in humans and humanized mice and is a promising target for new drug development. *Signal Transduct Target Ther* 6:1-12.

27) Morse JS, Lalonde T, Xu S, Liu WR (2020) Learning from the past: Possible urgent prevention and treatment options for severe acute respiratory infections caused by 2019-nCoV. *Chembiochem* 21:730-738.

28) Costanzi E, Kuzikov M, Esposito F, Albani S, Demitri N, et al. (2021) Structural and biochemical analysis of the dual inhibition of MG-132 against SARS-CoV-2 main protease (M^{pro} or $3CL^{pro}$) and human cathepsin-L. *Int J Mol Sci* 22:11779.

30) Santos R, Ursu O, Gaulton A, Bento AP, Donadi RS, et al. (2017) A comprehensive map of molecular drug targets. *Nat Rev Drug Discov* 16:19-34.

31) Wu S, Cao J, Wang T, Zhou M (2015) Experimental study on the antifungal and antiviral effects of volatile oil from artemisia argyi leaves on herpes zoster virus. *J Chi med* 34:70-71.

32) Han Y, Dai C, Tang L (2005) Preliminary study on the antiviral effect of volatile oil from Artemisia argyi leaves. *Amino Acids & Biotic Resources* 2:14-16.

34) Zhao Z, Wang L, Zheng L, Geng N, Zhao X, et al. (2015) The inhibitory effect of volatile oil from Artemisia argyi leaves on HBV. *Journal of Zhengzhou University* 50:301-304.

35) Ke M (1982) Physicochemical and pharmacological properties of effective ingredients in Chinese herbal medicine. Hunan Science and Technology Press.

36) Hai C, Bai X, Chen H, Yang X, Yang J, et al. (2021) Study on characteristic volatile components of chinese white Chrysanthemum by gc-ms combined with chemometrics. *Chemical Reagents* 43:584-589.

37) Ren A, Wang Z, Lu Z, Wang L, Wu Y (1999) Experimental study on the antibacterial and antiviral effects of wild Chrysanthemum. *Pharmaceutical Biotechnology* 4:241-244.

38) Zhao H, Zhao Y (2010) Study on effect of ethanol extracts from *rollius chinensis* bge. on resisting influenza virus *in vitro*. *China Pharmaceuticals* 1:10-11.

39) Guo X, Zhang L (2017) Comparative study on volatile oil components in *Chrysanthemum morifolium* from different producing areas. *Modern Chinese Medicine* 19:821-827.

40) Zhang X, Chen X, Wu Y (2021) Research progress on chemical constituents and pharmacological activities of volatile oil of aiye (Artemisia argyi). *Tradit Chin Med* 39:111-118.

41) Li JianJun LJ, Ren MeiLing RM, Shang XingChen SX, Lian XiaoYa LX, Wang HongLei WH (2017) Extraction of volatile oil from honeysuckle by distillation and its component analysis. *J Henan Agri Sci* 46:144-148.

42) Wang JianLing WJ, Ni KePing NK, Ji XiaoMing JX, Li BingJie LB, LüQuanJian L (2012) Analysis of the essential oil from *Trollius chinensis* bunge bygc-ms and its application in unblended cigarette flavoring. *J Henan Agri Sci* 41:121-124.

43) He S, Zhang C, Zhou P, Zhang X, Ye T, et al. (2019) Herb-induced liver injury: Phylogenetic relationship, structure-toxicity relationship, and herb-ingredient network analysis. *Int J Mol Sci* 20:3633.

44) Guo Y, Bao C, Ma D, Cao Y, Li Y, et al. (2019). Network-based combinatorial CRISPR-Cas9 screens identify synergistic modules in human cells. *ACS synthetic biology* 8:482-490.